

2

## REPORT DOCUMENTATION PAGE

1a NAME OF PERFORMING ORGANIZATION N/A			1b RESTRICTIVE MARKINGS N/A		
2a NAME OF PERFORMING ORGANIZATION University of Washington			3 DISTRIBUTION AVAILABILITY OF REPORT Distribution Unlimited		
4 PERFORMING ORGANIZATION REPORT NUMBER N/A			5 MONITORING ORGANIZATION REPORT NUMBER(S) N/A		
6a ADDRESS (City, State, and ZIP Code) Department of Chemistry BG-10 University of Washington Seattle, Washington 98195			7a NAME OF MONITORING ORGANIZATION Office of Naval Research		
8a NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research			9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-88-K-0201		
8b OFFICE SYMBOL (if applicable) ONR			10 SOURCE OF FUNDING NUMBERS		
8c ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, Virginia 22217-5000			PROGRAM ELEMENT NO 61153N		PROJECT NO PR04106
			TASK NO 44n003		WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) (U) Control of Synthetic Peptide Tertiary Structure					
12 PERSONAL AUTHOR(S) Paul B. Hopkins					
13a TYPE OF REPORT Annual		13b TIME COVERED FROM 2/89 TO 4/90		14 DATE OF REPORT (Year, Month, Day) 1990, April 15	
15 PAGE COUNT					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Peptide secondary structure, molecular design, circular dichroism, $\alpha$ -helix, protein thermostabilization, etc.		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) The proposed research seeks to design and synthesize large organic molecules possessing precisely defined geometric shapes, a capability which will make possible the design of catalysts and receptors. The approach entails the design of unnatural amino acids which, when incorporated into peptides, will enforce tertiary structure on the overall molecule. Methods are also suggested whereby transiently stable noncovalent complexes of peptides might be permanently stabilized by covalent crosslinking, leading to the concept of "self-assembly" of large, highly ordered organic molecules. The additional possibility of assembling large structures from smaller pieces by incorporation of temporary functionality which will initially spatially orient small pieces prior to covalent assembly is suggested, a concept termed "directed assembly."					
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION (U)		
22a NAME OF RESPONSIBLE INDIVIDUAL M. Marron			22b TELEPHONE (Include Area Code) (206) 696-4760		22c OFFICE SYMBOL ONR

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.

SECURITY CLASSIFICATION OF THIS PAGE

## DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

All other editions are obsolete.

## Annual Progress Report on Contract N00014-88-K-0201

**Principal Investigator:** Paul B. Hopkins  
**Contractor:** University of Washington  
**Contract Title:** Control of Synthetic Peptide Tertiary...  
**Start Date:** 1 February 1988  
**Current Date:** 15 April, 1990

### Research Objective:

The design, synthesis, and structural characterization of peptides of 10 to 35 residues with stable secondary and tertiary structure

### Progress (Year 2):

As described in the workplan of last year's annual report, we focused our attention this past year on the synthesis and evaluation of a series of peptides which contain two metal-ligating sides chains spaced such that simultaneous chelation of a single metal ion might enhance  $\alpha$ -helicity. We chose as our goal the synthesis of eight peptides,  $\text{AcAdaAla}_m\text{Ada}(\text{Ala}_4\text{GluLys})_3\text{NH}_2$ , in which Ada is an L- $\alpha$ -amino acid residue with the aminodiacetic acid side chain,  $(\text{CH}_2)_n\text{N}(\text{CH}_2\text{CO}_2\text{H})_2$ , and values of m and n are all combinations of  $m=2,3$  and  $n=1-4$ . Half of these syntheses are now complete. The syntheses require, at the outset, the availability of the four L- $\alpha$ -amino acids represented by Ada. These compounds were synthesized as the N $^\epsilon$ -Boc-Ada(O $^\epsilon$  Bzl) $_2$ -protected derivatives and were incorporated into peptides by solid phase methods in either a fully stepwise synthesis, or, in a few cases, by a block approach. The resulting peptides were purified by reverse phase HPLC. FABS MS analysis of the purified peptides afforded the expected parent ion; for one peptide (prepared without terminal N-acetylation) sequence analysis confirmed the expected structure.

Circular dichroism analysis of these synthetic peptides is currently in progress. The limited available information suggests that the tether length (m) and spacing between ligating residues (n) are critical. As expected, long tethers apparently decouple the ligation of the metal from significant effect on the conformation of the attached peptide. The impact of the shortest tethers is at present unknown. Limited data also suggest that the spacing between ligating residues is critical, with  $n=2$  being apparently too short to allow helix formation and actually destabilizing the  $\alpha$ -helical form. The

equilibrium ratio of helix to random coil was increased three fold by addition of  $\text{Cu}^{2+}$  to one of these peptides. Whether this is the greatest enhancement for the family of eight peptides will be known only on completion of the syntheses and further circular dichroism analysis.

### **Workplan (Year 3):**

Having (at last!) prepared a family of peptides which exhibit metal ion-dependent conformational properties, the focus of the project will now shift to evaluation of the solution conformational properties of these peptides. We will determine the CD spectra from 180 to 260 nm in the presence of a wide variety of metal ions ( $\text{Mg}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Co}^{2+}$ ). Theory predicts that a maximum shift of about  $10^5$  of the equilibrium constant in favor of  $\alpha$ -helix could be observed; of the three peptides we have thus far evaluated we have seen at most a factor of 3. Based upon those values of  $m$  and  $n$  (see above) which are found to provide the most enhanced helicity, we may elect to synthesize one or two more peptides in which each peptide bears two different values of  $n$ . The goal of this would be to optimize the stabilization of the helical form. We might, for example, prepare peptides in which the first ligand is held with an ethylene and the second with a propylene tether and vice versa. For those combinations of peptide/metal which are found to be interesting by CD, we will study the impact of pH on the conformational equilibria. The state of aggregation of these peptides in their helical form (expected to be monomers) will be verified by gel filtration experiments. Nuclear magnetic resonance studies will be undertaken to verify the helical structure evident by CD. Finally, computer molecular modeling studies, using molecular mechanics techniques, will be undertaken to attempt to rationalize the observed trends in helix stabilization as a function of tether length and inter-residue spacing.

### **Inventions:**

None

### **Publications:**

Hopkins, Paul B. Hopkins, Ruan, F.; Chen, Y.; Sasaki, T.; and Itoh, K., "Metal Ion-Induced  $\alpha$ -Helicity in Synthetic Peptides Containing Unnatural, Metal-Ligating Residues," Abstract (submitted) 200th National Meeting of the American Chemical Society, Washington D.C., August 26-31, 1990.

As soon as circular dichroism studies are complete on the eight peptides described above, we plan to submit one or more manuscripts to the *Journal of the American Chemical Society*.

### Training Activities:

Mr. Fugiang Ruan and Dr. Yanqiu Chen, both native born citizens of the Peoples Republic of China are currently working on this project. Mr. Ruan's major focus is peptide synthesis; Dr. Chen is evaluating the solution conformational properties of the synthetic peptides. Professor Tomikazu Sasaki, a new faculty member with considerable experience in the field of peptide design and synthesis, is serving as an informal consultant on the project.

### Awards/Fellowships:

The PI is an Alfred P. Sloan Fellow (1988-1992) and is recipient of an NIH Research Career Development Award. Fuqiang Ruan is a PRC Scholar.

Accession For	
NEWS GRA&I	<input checked="" type="checkbox"/>
IC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	